



Discharge Advice on Electrophysiology & Radiofrequency Ablation

How do I take care of my wound?



Caring for wound

- Inspects wound site(s) by looking in the mirror to check for any signs and symptoms of wound infection or bleeding.
- Always keep your wound site clean and dry, especially at affected groin(s) area.

Recognising the signs of infection:

- A sudden increase in pain over the wound site.
- Increased swelling, redness and heat felt around the wound.
- ♥ Increased amount of discharge or presence of pus-like drainage.
- ▼ Fever of more than 38°.



Recognising the signs of wound bleeding:

- Active bleeding at wound site.
- Swelling at affected groin area and/or thigh.
- Increasing size of bruise.
- Worsening pain at wound area.
- ♥ Giddiness or syncope (loss of consciousness).
- Increasing fatigue.
- Chest pain or discomfort.
- Shortness of breath.

When should I see a doctor?



Follow-up appointments

- You are recommended to attend all follow-up appointments as scheduled.
- Please call our appointment line at 6908 2222 if you did not receive your appointment details within 2-3 days after discharge.
- Please call the appointment line 5 days prior to the scheduled appointment if you are unable to attend the session.
- Do bring along your appointment card any relevant memo(s), discharge summary and test results from your doctor(s) whom you have consulted outside of NUH, if any.
- You may be expected to have an electrocardiogram performed at least 30 minutes prior to your scheduled follow-up appointment time.



Importance of follow-up appointments with your Electrophysiology Cardiologist:

♥ Wound assessments and further interventions/investigations may be carried out during your follow-up appointments that are scheduled from 6-12 weeks after your procedure, to ensure maintenance of the therapeutic outcomes.

When to seek immediate medical attention?

- New, unexplained heart symptoms and/or similar signs and symptoms that you had experienced before your treatment/procedure.
- Signs and symptoms of wound infection.
- Signs and symptoms of wound bleeding.
- Worsening numbness or weakness at affected limb(s).
- Giddiness, syncope, unexplained lethargy or shortness of breath.






Where should I seek immediate medical attention?

 Emergency Department at the National University Hospital (NUH) or any hospital nearest to you.

Important information to note

- ♥ You can resume your daily activities within 24 hours after your procedure with few restrictions. These include:
 - Light activities for the first 2 weeks.
 - Avoid strenuous activities that may strain your wound for 1 week after the procedure. These include activities that:
 - Involve bending over 90° which may affect your wound at the groin(s) area, such as squatting.
 - Involve the use of core muscles that affect the abdominal and groin(s) area such as sit-ups or abdominal crunches.
- When in doubt, please do not hesistate to contact us at 6908 2222.
- Further instructions:



National University Heart Centre, Singapore (NUHCS)

NUHCS Heart Clinic @ National University Hospital (NUH)

NUH Main Building Zone F

Opening Hours: 8.00 am - 5.30 pm (Monday - Friday)

Closed on Weekends & Public Holidays

OneNUHS Hotline: (65) 6908 2222

OneNUHS Appointments: appointment@nuhs.edu.sg
OneNUHS General Enquiries: contactus@nuhs.edu.sg

www.nuhcs.com.sg



Scan for more information

Take charge of your health and access health information and services across NUHS institutions.

Download the OneNUHS App now!





Scan to download



Every day, we impact lives by providing assistance to financially disadvantaged patients.





Scan here to donate

The information in this brochure is meant for educational purposes and should not be used as a substitute for medical diagnosis or treatment. Please seek your doctor's advice before starting any treatment, or if you have any questions related to your health, physical fitness or medical condition.

© 2023, National University Hospital (Singapore) Pte. Ltd. All rights reserved. No part of this publication may be reproduced or shared without prior permission from National University Hospital (Singapore) Pte. Ltd.

Information is correct at time of printing (Jun 2023) and subject to revision without prior notice.